

COMPARISON OF TREATMENTS FOR URINARY INCONTINENCE:

GENITYTE®	NON-SURGICAL PROCEDURE	RISKS	RECOVERY	SUCCESS RATE WITH GENITYTE
<p>① Stress Urinary Incontinence</p> <p>② Mixed Urinary Incontinence (Stress & Urge)</p> <p>③ Vaginal Wall Prolapse</p> <p>④ Rectal Incontinence</p>	<p><u>Non-Invasive</u></p> <p><u>Uses Infrared Light</u> Stimulates fibroblasts to repair skin, muscle and connective tissue.</p> <p><u>Tones and Strengthens the Pelvic Floor</u> Tones the urethra and its supporting tissues, vaginal walls, perineum, anus and pelvic floor muscles.</p>	<p><u>Worsening of Urinary Urge</u> -7 out of 345 procedures -Temporary: due to dynamic re-positioning of the urethra -Resolved within 2-4 weeks</p>	<p><u>No Anesthesia</u></p> <p><u>No Pain</u></p> <p><u>No Recovery</u></p>	<p><u>Stress Incontinence Treatment:</u> -All patients report at least 80% improvement, 7/10 report complete resolution of leaking</p> <p><u>Mixed Incontinence Treatment:</u> -All patients report at least 70% improvement, 4/10 report complete resolution of leak/urge</p> <p><u>Vaginal Wall Prolapse Treatment:</u> -All patients have vaginal wall prolapse improvement -Some have complete resolution</p> <p><u>Rectal Incontinence treatment with Genityte:</u> -All patients have rectal tone improvement -Majority of patients have resolution of rectal incontinence</p>
"THE SLING" SURGERY	NON-SURGICAL PROCEDURE	RISKS	RECOVERY	SUCCESS RATE
<p>① Stress Urinary Incontinence</p>	<p><u>Invasive</u> -Foreign material is surgically placed in the tissues under the urethra</p> <p><u>Uses Foreign Material</u> -Sling material, usually 'mesh' is secured with sutures -Positioned like a hammock under the urethra</p> <p><u>Tones and Strengthens the tissue under the urethra and above the vaginal wall</u></p>	<p><u>Erosion of Sling Material</u> -Through the Vaginal Wall -Through the Urethra</p> <p><u>Urethral Obstruction</u> -Unable to Start Urination</p> <p><u>Urinary Retention</u> -Unable to Empty Bladder</p> <p><u>De Novo Urge Incontinence</u> -New Urge Incontinence</p> <p><u>Severe Bleeding, Infection</u></p>	<p><u>General Anesthesia</u></p> <p><u>Spinal Anesthesia</u></p> <p><u>4-6 weeks Recovery</u></p> <p><u>Activity Restrictions</u> -High impact exercise restriction</p> <p><u>Lifting Restrictions</u> -Lifetime restriction may not lift more than 20#</p>	<p><u>Stress Incontinence treatment with the Sling Surgery:</u> -Up to 89% of patients have significant improvement -However; complication rates have been reported as high as 31%</p>

